# DEPARTMENT OF THE ARMY HEADQUARTERS, UNITED STATES ARMY MEDICAL COMMAND 2050 Worth Road Fort Sam Houston, Texas 78234-6000

MEDCOM Circular No. 40-12

1 August 2003

## Expires 1 August 2005 Medical Services TOBACCO CESSATION OUTPATIENT FORM

**1. HISTORY.** This issue publishes a revision of this publication.

#### 2. PURPOSE.

- a. This circular provides policy and implementing instructions for use of the tobacco cessation outpatient form prescribed by this circular: U.S. Army Medical Command (MEDCOM) Form 709-R (Tobacco Cessation Documentation).
- b. This form will facilitate outpatient treatment record (OTR) documentation by cueing practitioners to document key aspects in their assessment and treatment of patients who use tobacco products. A panel of expert consultants from the Army, Navy, Air Force, and Department of Veterans Affairs (VA) identified key aspects by thoroughly examining scientific evidence on tobacco use cessation. This panel synthesized the evidence on promoting tobacco use cessation in the Department of Defense/VA Clinical Practice Guideline to promote tobacco use cessation in the primary care setting. Key aspects were then transformed onto the form named in paragraph a above and prescribed by this circular. Use of this form is not mandatory, rather it is designed to facilitate documentation of care provided to patients who are attempting tobacco use cessation in primary care.
- **3. APPLICABILITY**. This circular applies to all MEDCOM facilities. All facilities are granted use of the test form (prescribed herein) to document care of patients who use tobacco products.
- **4. REFERENCES**. AR 40-66, Medical Record Administration and Health Care Documentation, provides guidance on medical record documentation and is applicable.

<sup>\*</sup>This circular supersedes MEDCOM Circular 40-12, 1 August 2001, including change.

#### 5. EXPLANATION OF ABBREVIATIONS AND TERMS.

a. Abbreviations.

MEDCOM......U.S. Army Medical Command OTR.....outpatient treatment record SF.....standard form VA......Department of Veterans Affairs

- b. Terms. See AR 40-66.
- 6. RESPONSIBILITIES. See AR 40-66.

#### 7. POLICY.

- a. Personnel in military treatment facilities may use MEDCOM Form 709-R for the period of the test (through 1 August 2005) or as directed by the MEDCOM.
- b. The MEDCOM test form prescribed by this circular will be filed in the OTR with the standard form (SF) 600 (Health Record-Chronological Record of Medical Care) in reverse chronological order (most recent on top).
- c. MEDCOM Form 709-R may be used in lieu of the SF 600 to document treatment only for tobacco use cessation patients being treated on an outpatient basis.
- d. All current requirements of AR 40-66, other than those addressed in this circular, remain in effect.
- **8. INSTRUCTIONS FOR USE OF THE TOBACCO CESSATION DOCUMENTATION FORM.** Note: This form is authorized for local reproduction (that is, "-R" form) and is contained in appendix A of this circular. This form is to be printed head to foot.

MEDCOM Form 709-R.

- a. Purpose. This form may be used to document the treatment to promote tobacco use cessation.
- b. Preparation. This form has five sections. Section I, vital signs, is to be completed by the technician. Section II, patient assessment, is to be completed by the patient and reviewed by the provider. Section III, medical history and physical assessment, is to be completed by the health care provider. Sections IV and V, assessment and action plan, are to be completed by the health care provider.
- c. Content. Section I includes documentation of height, weight, and vital signs. Section II includes questions on tobacco use history. Section III includes check box and

free-hand areas for documentation of the patient's medical history and physical assessment. Section IV is the primary assessment and Section V is an action plan.

### APPENDIX A

Appendix A contains the following "-R" form (authorized for local reproduction).

MEDCOM Form 709-R (Tobacco Cessation Documentation)

| MEDICAL RECORD - TOBACCO CESSATION DO  For use of this form see MEDCOM Circular 4  |             | NTAT    | ION      | TREATMENT FAC          | ILITY D       | ATE              |
|--|-------------|---------|----------|------------------------|---------------|------------------|
| SECTION 1 - VITA   | L SIGNS     | (Comp   | leted by | Technician)            |               |                  |
| TIME: BP: PULSE:   | RESP        | :       |          | TEMP:                  | HT:           | WT:              |
| ALLERGY: ME  | -<br>DICATI | ONS:    |          |                        |               | <del></del>      |
| SECTION II - PATIENT ASSESSM   | AENT (Co    | mplete  | d by Pat | ient/reviewed by Provi | der)          |                  |
| At what age did you start using tobacco?   |             |         |          |                        |               |                  |
| 2. What type(s) and amount(s) of tobacco do you  | use?        |         |          |                        |               |                  |
| TYPE(S)  | YES         | NO      |          | AMOUNT(S)              | PER DA        | Y PER MONTH      |
| a. Cigarette   |             |         | Pack     | s                      |               |                  |
| b. Pipe  |             |         | Bowl     | S                      |               |                  |
| c. Cigar   |             |         | Cigar    | 'S                     |               |                  |
| d. Snuff   |             |         | Cans     |                        |               |                  |
| e. Chew  |             |         | Pouc     | h                      |               |                  |
| 3. How soon after you wake up do you use tobac   | co?         |         | After 3  | 80 minutes 🔲 W         | ithin 30 m    | ninutes          |
| 4. Have you quit before?   |             |         |          |                        |               |                  |
| 5. How many times have you quit before?  |             |         |          |                        |               |                  |
| 6. What was the longest period you were able to  | quit?       |         |          |                        |               |                  |
| 7. What caused you to start using tobacco again?   |             |         |          |                        |               |                  |
|  |             |         |          |                        |               |                  |
|  |             |         |          |                        |               |                  |
|  |             |         |          |                        |               |                  |
| 8. Did you use any of the following to help you qu   | uit?        | F       | Patch    | Gum Z                  | yban 🗌        | Inhaler          |
| ☐ Nasal Spray ☐ Individual Counseling ☐  | ] Forma     | al Pro  | gram     | Other                  | 7,000         |                  |
| 9. Why do you want to quit tobacco use?  | Financi     | al Sa   | ving     | ☐ Breathing Pro        | blems [       | ] Heart Problems |
| Fear of Cancer Family/Social Pressure  | е 🗌         | Other   | Issues   |                        |               |                  |
| 10. What support do you have available to help yo  | ou quit 1   | tobac   | co use?  | Family                 |               | ] Friends        |
| ☐ Work ☐ Other   |             |         |          |                        |               | ,<br>-           |
| 11. What type of program do you believe would h  | nelp you    | ı the   | most?    |                        |               |                  |
|  | Counse      |         |          | Self Quit              |               |                  |
|  |             |         |          |                        |               |                  |
| PATIENT'S IDENTIFICATION (For typed or written entries give<br>first, middle; grade; date; hospital or medical facility) | e: Name     | - last, |          |                        |               |                  |
|  |             |         |          |                        |               |                  |
|  |             |         | İ        | **                     |               |                  |
|  |             |         |          |                        |               |                  |
|  |             |         |          |                        |               |                  |
|  |             |         |          | (Patient's             | s Signature/l | Date)            |

|  | PHYSICAL ASSESSMENT (Completed by Health Care Provider)   | 48.4         |
|--|---|--------------|
| MEDICATIONS REVIEWED: ☐ Yes ☐ No ALLI  | IEDICAL HISTORY ERGIES REVIEWED: Yes No LMP:  |              |
| •  |   |              |
|  |   |              |
| ETOH: To Vas To No. To Cut Down.   | Annoyed   |              |
| During the past month have you been bothered by  |   | ] No         |
|  | Little interest or pleasure in doing things  Yes [  | No           |
| PMH affecting use of NRT/Bupropion:  |   |              |
| PRECAUTIONS/CONTRAINDICATIONS YES  | S NO PRECAUTIONS/CONTRAINDICATIONS YES MOOD DISORDER  | NO           |
| SEIZURES   | POLYCYTHEMIA  | <del> </del> |
| CHRONIC PAIN DISORDER  | DIABETES  |              |
| LIVER DISEASE  | CARDIOVASCULAR DISEASE  |              |
| HYPERTHYROIDISM  | COMPLICATIONS OF TOBACCO USE YES  | NO           |
| KIDNEY DISEASE PREGNANCY   | CHRONIC OBSTRUCTIVE PULMONARY DZ ASTHMA   | +            |
| LACTATING  | CORONARY ARTERY DISEASE   | 1            |
| SUBSTANCE ABUSE  | CANCER  |              |
| EATING DISORDER  | ERECTILE DYSFUNCTION  |              |
| POST TRAUMATIC STRESS DISORDER   | PERIPHERAL VASCULAR DISEASE OTHER   | -            |
| ANXIETY Physical Assessment:   | OTHER   |              |
|  |   |              |
|  |   |              |
|  | MENT (Completed by Health Care Provider)  | pengananan   |
| PRIMARY ASSESSMENT: Tobacco Cessation V6   | 5.49 4 (DOD unique extender) ICD - 9-CM 305.1   |              |
| PRIMARY ASSESSMENT: Tobacco Cessation V6 SECTION V - ACTION  | 5.49 4 (DOD unique extender) ICD - 9-CM 305.1  PLAN (Completed by Health Care Provider)   | All says     |
| PRIMARY ASSESSMENT: Tobacco Cessation V6: SECTION V - ACTION  1. MEDICATIONS: NRT Prescribed?  | 5.49 4 (DOD unique extender) ICD - 9-CM 305.1  PLAN (Completed by Health Care Provider)  ES NO  |              |
| PRIMARY ASSESSMENT: Tobacco Cessation V6  SECTION V - ACTION  1. MEDICATIONS: NRT Prescribed?  YI  Transdermal Nicotine (Contraindicated in P  | 5.49 4 (DOD unique extender) ICD - 9-CM 305.1  PLAN (Completed by Health Care Provider)  ES  NO  Pregnancy)  7 Mg x weeks  14 Mg x  |              |
| PRIMARY ASSESSMENT: Tobacco Cessation V6:  SECTION V - ACTION  1. MEDICATIONS: NRT Prescribed?  YI  Transdermal Nicotine (Contraindicated in P  21 Mg x weeks  Mg x  | 5.49 4 (DOD unique extender) ICD - 9-CM 305.1  PLAN (Completed by Health Care Provider)  ES  NO  Pregnancy)  7 Mg x weeks  14 Mg x  |              |
| PRIMARY ASSESSMENT: Tobacco Cessation V6  SECTION V - ACTION  1. MEDICATIONS: NRT Prescribed?  YI  Transdermal Nicotine (Contraindicated in P  21 Mg x weeks  Mg x Polacrilex Nicotine PRN   | 5.49 4 (DOD unique extender) ICD - 9-CM 305.1  PLAN (Completed by Health Care Provider)  ES  NO  Pregnancy)  7 Mg x weeks  14 Mg x weeks  |              |
| PRIMARY ASSESSMENT: Tobacco Cessation V6:  SECTION V - ACTION  1. MEDICATIONS: NRT Prescribed?  YI  Transdermal Nicotine (Contraindicated in P  21 Mg x weeks  Mg x  | 5.49 4 (DOD unique extender) ICD - 9-CM 305.1  PLAN (Completed by Health Care Provider)  ES  NO  Pregnancy)  7 Mg x weeks  14 Mg x weeks  |              |
| PRIMARY ASSESSMENT: Tobacco Cessation V6  SECTION V - ACTION  1. MEDICATIONS: NRT Prescribed?  YI  Transdermal Nicotine (Contraindicated in P  21 Mg x weeks  Mg x Polacrilex Nicotine PRN   | 5.49 4 (DOD unique extender) ICD - 9-CM 305.1  PLAN (Completed by Health Care Provider)  ES  NO  Pregnancy)  7 Mg x weeks  14 Mg x weeks  |              |
| PRIMARY ASSESSMENT: Tobacco Cessation V6  SECTION V - ACTION  1. MEDICATIONS: NRT Prescribed? YI  Transdermal Nicotine (Contraindicated in P  21 Mg x weeks Mg x Polacrilex Nicotine PRN  Other: Bupropion SR 150 mg   | 5.49 4 (DOD unique extender) ICD - 9-CM 305.1  PLAN (Completed by Health Care Provider)  ES NO  Pregnancy) 7 Mg x weeks 14 Mg x weeks  po,qd x days, then bid.  |              |
| PRIMARY ASSESSMENT: Tobacco Cessation V6  SECTION V - ACTION  1. MEDICATIONS: NRT Prescribed? YI  Transdermal Nicotine (Contraindicated in P  21 Mg x weeks Mg x Polacrilex Nicotine PRN  Other: Bupropion SR 150 mg  2. Tobacco Cessation Counseling:   | 5.49 4 (DOD unique extender) ICD - 9-CM 305.1  PLAN (Completed by Health Care Provider)  ES   |              |
| PRIMARY ASSESSMENT: Tobacco Cessation V6  SECTION V - ACTION  1. MEDICATIONS: NRT Prescribed? YI  Transdermal Nicotine (Contraindicated in P  21 Mg x weeks Mg x  Polacrilex Nicotine PRN  Other: Bupropion SR 150 mg  2. Tobacco Cessation Counseling:  Patient congratulated on decision to quit to  | 5.49 4 (DOD unique extender) ICD - 9-CM 305.1  PLAN (Completed by Health Care Provider)  ES   |              |
| PRIMARY ASSESSMENT: Tobacco Cessation V6  SECTION V - ACTION  1. MEDICATIONS: NRT Prescribed? YI  Transdermal Nicotine (Contraindicated in P  21 Mg x weeks  Mg x  Polacrilex Nicotine PRN  Other: Bupropion SR 150 mg  2. Tobacco Cessation Counseling: Patient congratulated on decision to quit to patient advised to avoid all tobacco productions.  | 5.49 4 (DOD unique extender) ICD - 9-CM 305.1  PLAN (Completed by Health Care Provider)  ES   |              |
| PRIMARY ASSESSMENT: Tobacco Cessation V6  SECTION V - ACTION  1. MEDICATIONS: NRT Prescribed? YI  Transdermal Nicotine (Contraindicated in Parameter Paramet | 5.49 4 (DOD unique extender)  PLAN (Completed by Health Care Provider)  ES NO  Pregnancy) 7 Mg x weeks 14 Mg x pounds  weeks  po,qd x days, then bid.  obacco usage: Quit Date  cts during NRT.   |              |
| PRIMARY ASSESSMENT: Tobacco Cessation V6  SECTION V - ACTION  1. MEDICATIONS: NRT Prescribed? YI  Transdermal Nicotine (Contraindicated in Parameter Paramet | 5.49 4 (DOD unique extender)  PLAN (Completed by Health Care Provider)  ES  |              |
| PRIMARY ASSESSMENT: Tobacco Cessation V6  SECTION V - ACTION  1. MEDICATIONS: NRT Prescribed? YI  Transdermal Nicotine (Contraindicated in P  21 Mg x weeks  Mg x  Polacrilex Nicotine PRN  Other: Bupropion SR 150 mg  2. Tobacco Cessation Counseling:  Patient congratulated on decision to quit to  Patient advised to avoid all tobacco product  Tobacco cessation benefits reviewed.  Patient advised of withdrawal symptoms.  Patient concerns and support systems add  | 5.49 4 (DOD unique extender)  PLAN (Completed by Health Care Provider)  ES  |              |
| PRIMARY ASSESSMENT: Tobacco Cessation V6  SECTION V - ACTION  1. MEDICATIONS: NRT Prescribed? YI  Transdermal Nicotine (Contraindicated in P  21 Mg x weeks  Mg x Polacrilex Nicotine PRN  Other: Bupropion SR 150 mg  2. Tobacco Cessation Counseling: Patient congratulated on decision to quit to Patient advised to avoid all tobacco product Patient advised of withdrawal symptoms. Patient concerns and support systems add Patient advised to take medication as direct  | FLAN (Completed by Health Care Provider)  ES NO  Pregnancy) 7 Mg x weeks 14 Mg x weeks po,qd x days, then bid.  Obacco usage: Quit Date cts during NRT.   |              |
| PRIMARY ASSESSMENT: Tobacco Cessation V6  SECTION V - ACTION  1. MEDICATIONS: NRT Prescribed? YI  Transdermal Nicotine (Contraindicated in P  21 Mg x weeks Mg x Polacrilex Nicotine PRN  Other: Bupropion SR 150 mg  2. Tobacco Cessation Counseling: Patient congratulated on decision to quit to Patient advised to avoid all tobacco production in Patient advised of withdrawal symptoms. Patient advised of withdrawal symptoms. Patient concerns and support systems add Patient advised to take medication as direction in Educational materials given to patient.  What type of tobacco cessation program would be producted in Patient.  | FLAN (Completed by Health Care Provider)  ES NO  Pregnancy) 7 Mg x weeks 14 Mg x weeks po,qd x days, then bid.  Obacco usage: Quit Date cts during NRT.   | weeks        |
| PRIMARY ASSESSMENT: Tobacco Cessation V6  SECTION V - ACTION  1. MEDICATIONS: NRT Prescribed? YI  Transdermal Nicotine (Contraindicated in P  21 Mg x weeks Mg x Polacrilex Nicotine PRN  Other: Bupropion SR 150 mg  2. Tobacco Cessation Counseling: Patient congratulated on decision to quit to Patient advised to avoid all tobacco product Patient advised of withdrawal symptoms. Patient advised of withdrawal symptoms. Patient concerns and support systems add Patient advised to take medication as direct Educational materials given to patient.  What type of tobacco cessation program would be producted to the program would be producted to the patient.  | PLAN (Completed by Health Care Provider)  ES NO Pregnancy) 7 Mg x weeks 14 Mg x po,qd x days, then bid.  Obacco usage: Quit Date cts during NRT.  | weeks        |
| PRIMARY ASSESSMENT: Tobacco Cessation V6  SECTION V - ACTION  1. MEDICATIONS: NRT Prescribed? YI  Transdermal Nicotine (Contraindicated in P  21 Mg x weeks Mg x Polacrilex Nicotine PRN  Other: Bupropion SR 150 mg  2. Tobacco Cessation Counseling: Patient congratulated on decision to quit to Patient advised to avoid all tobacco product Patient advised of withdrawal symptoms. Patient advised of withdrawal symptoms. Patient concerns and support systems add Patient advised to take medication as direct Educational materials given to patient.  What type of tobacco cessation program wout Formal Group Behavior  | PLAN (Completed by Health Care Provider)  ES NO Pregnancy) 7 Mg x weeks 14 Mg x po,qd x days, then bid.  Obacco usage: Quit Date cts during NRT.  | weeks        |
| PRIMARY ASSESSMENT: Tobacco Cessation V6  SECTION V - ACTION  1. MEDICATIONS: NRT Prescribed? YI  Transdermal Nicotine (Contraindicated in P  21 Mg x weeks Mg x Polacrilex Nicotine PRN  Other: Bupropion SR 150 mg  2. Tobacco Cessation Counseling: Patient congratulated on decision to quit to Patient advised to avoid all tobacco product Patient advised of withdrawal symptoms. Patient advised of withdrawal symptoms. Patient advised to take medication as direct Educational materials given to patient.  3. What type of tobacco cessation program wout Formal Group Behavior  4. Referral To:     Stress Management Dietary   | PLAN (Completed by Health Care Provider)  ES NO  Pregnancy) 7 Mg x weeks 14 Mg x  weeks  po,qd x days, then bid.  Obacco usage: Quit Date cts during NRT.  Iressed.  Id you like to participate in?  Modification 0 One On One Self Quit Prog | weeks        |
| PRIMARY ASSESSMENT: Tobacco Cessation V6  SECTION V - ACTION  1. MEDICATIONS: NRT Prescribed? YI  Transdermal Nicotine (Contraindicated in P  21 Mg x weeks Mg x Polacrilex Nicotine PRN  Other: Bupropion SR 150 mg  2. Tobacco Cessation Counseling: Patient congratulated on decision to quit to Patient advised to avoid all tobacco production Tobacco cessation benefits reviewed. Patient advised of withdrawal symptoms. Patient advised of withdrawal symptoms. Patient advised to take medication as direction and the Patient advised to take medication as direction and the Patient advised to take medication as direction and the Patient advised to take medication as direction and the Patient advised to take medication as direction and the Patient advised to take medication as direction and the Patient advised to take medication as direction and the Patient advised to take medication as direction and the Patient advised to take medication as direction and the Patient advised to take medication as direction and the Patient advised to take medication as direction and the Patient advised to take medication as direction and the Patient advised to take medication as direction and the Patient advised to take medication as direction and the Patient advised to take medication as direction and the Patient advised to take medication as direction and the Patient advised to take medication as direction and the Patient advised to take medication as direction and the Patient advised to take medication and the Patient advised to take medication as direction and the Patient advised to take medication as direction and the Patient advised to take medication as direction and the Patient advised to take medication and the Pati                       | PLAN (Completed by Health Care Provider)  ES NO  Pregnancy) 7 Mg x weeks 14 Mg x  weeks  po,qd x days, then bid.  Obacco usage: Quit Date cts during NRT.  Iressed.  Id you like to participate in?  Modification 0 One On One Self Quit Prog | weeks        |

The proponent of this publication is the Quality Management Directorate. Users are invited to send comments and suggested improvements on DA Form 2028 (Recommended Changes to Publications and Blank Forms) to Commander, U.S. Army Medical Command, ATTN: MCHO-Q, 2050 Worth Road, Fort Sam Houston, TX 78234-6026.

#### FOR THE COMMANDER:



KENNETH L. FARMER, JR. Major General Chief of Staff

EILEEN B. MALONE Colonel, AN Assistant Chief of Staff for Information Management

#### **DISTRIBUTION:**

This publication is available in electronic media only and is intended for MEDCOM distribution Ca and Da.

#### SPECIAL DISTRIBUTION:

MCHC (Stockroom) (1 cy)
MCHC (Library) (1 cy)
MCHS-AS (Forms Mgr) (1 cy)
MCHS-AS (Editor) (2 cy)